

ON THE WILKER AND HUYGENS-TYPE INEQUALITIES

CHAO-PING CHEN* AND RICHARD B. PARIS

ABSTRACT. Chen and Cheung [C.-P. Chen, W.-S. Cheung, Sharpness of Wilker and Huygens type inequalities, J. Inequal. Appl. 2012 (2012) 72, <http://dx.doi.org/10.1186/1029-242X-2012-72>] established sharp Wilker and Huygens-type inequalities. These authors also proposed three conjectures on Wilker and Huygens-type inequalities. In this paper, we consider these conjectures. We also present sharp Wilker and Huygens-type inequalities.

1. Introduction

Wilker [18] proposed the following two open problems:

(a) Prove that if $0 < x < \pi/2$, then

$$\left(\frac{\sin x}{x}\right)^2 + \frac{\tan x}{x} > 2. \quad (1.1)$$

(b) Find the largest constant c such that

$$\left(\frac{\sin x}{x}\right)^2 + \frac{\tan x}{x} > 2 + cx^3 \tan x$$

for $0 < x < \pi/2$. In [17], the inequality (1.1) was proved, and the following inequality

$$2 + \left(\frac{2}{\pi}\right)^4 x^3 \tan x < \left(\frac{\sin x}{x}\right)^2 + \frac{\tan x}{x} < 2 + \frac{8}{45} x^3 \tan x, \quad 0 < x < \frac{\pi}{2} \quad (1.2)$$

was also established, where the constants $(2/\pi)^4$ and $\frac{8}{45}$ are the best possible.

The Wilker-type inequalities (1.1) and (1.2) have attracted much interest of many mathematicians and have motivated a large number of research papers involving different proofs, various generalizations and improvements (cf. [1–3, 6, 8, 10–16, 19–23, 25–28] and the references cited therein).

A related inequality that is of interest to us is Huygens' inequality [9], which asserts that

$$2 \left(\frac{\sin x}{x}\right) + \frac{\tan x}{x} > 3, \quad 0 < |x| < \frac{\pi}{2}. \quad (1.3)$$

Wu and Srivastava [21, Lemma 3] established another inequality

$$\left(\frac{x}{\sin x}\right)^2 + \frac{x}{\tan x} > 2, \quad 0 < |x| < \frac{\pi}{2}. \quad (1.4)$$

Neuman and Sándor [15, Theorem 2.3] proved that for $0 < |x| < \pi/2$,

$$\frac{\sin x}{x} < \frac{2 + \cos x}{3} < \frac{1}{2} \left(\frac{x}{\sin x} + \cos x\right). \quad (1.5)$$

*Corresponding Author.

2010 Mathematics Subject Classification. 26D05.

Key words and phrases. Wilker and Huygens-type inequalities; Trigonometric functions; Bernoulli numbers.

By multiplying both sides of inequality (1.5) by $x/\sin x$, we obtain that for $0 < |x| < \pi/2$,

$$\frac{1}{2} \left[\left(\frac{x}{\sin x} \right)^2 + \frac{x}{\tan x} \right] > \frac{2(x/\sin x) + x/\tan x}{3} > 1. \quad (1.6)$$

Chen and Sándor [6] established the following inequality chain:

$$\begin{aligned} \frac{(\sin x/x)^2 + \tan x/x}{2} &> \left(\frac{\sin x}{x} \right)^2 \left(\frac{\tan x}{x} \right) > \frac{2(\sin x/x) + \tan x/x}{3} \\ &> \left(\frac{\sin x}{x} \right)^{2/3} \left(\frac{\tan x}{x} \right)^{1/3} > \frac{1}{2} \left[\left(\frac{x}{\sin x} \right)^2 + \frac{x}{\tan x} \right] > \frac{2(x/\sin x) + x/\tan x}{3} > 1 \end{aligned} \quad (1.7)$$

for $0 < |x| < \pi/2$.

In analogy with (1.2), Chen and Cheung [3] established sharp Wilker and Huygens-type inequalities. For example, these authors proved that for $0 < x < \pi/2$,

$$2 + \frac{8}{45}x^4 + \frac{16}{315}x^5 \tan x < \left(\frac{\sin x}{x} \right)^2 + \frac{\tan x}{x} < 2 + \frac{8}{45}x^4 + \left(\frac{2}{\pi} \right)^6 x^5 \tan x, \quad (1.8)$$

where the constants $\frac{16}{315}$ and $(2/\pi)^6$ are best possible,

$$\left(\frac{x}{\sin x} \right)^2 + \frac{x}{\tan x} < 2 + \frac{2}{45}x^3 \tan x, \quad (1.9)$$

where the constant $\frac{2}{45}$ is best possible, and

$$3 + \frac{3}{20}x^3 \tan x < 2 \left(\frac{\sin x}{x} \right) + \frac{\tan x}{x} < 3 + \left(\frac{2}{\pi} \right)^4 x^3 \tan x, \quad (1.10)$$

where the constants $\frac{3}{20}$ and $(2/\pi)^4$ are best possible.

In view of (1.8), (1.9) and (1.10), Chen and Cheung [3] posed the following conjectures.

Conjecture 1.1. For $0 < x < \pi/2$ and $n \geq 3$,

$$\begin{aligned} &2 + \sum_{k=3}^n \frac{(2(2^{2k}-1)|B_{2k}| - (-1)^k)2^{2k-1}}{(2k)!} x^{2k-2} \\ &+ \frac{(2(2^{2n+2}-1)|B_{2n+2}| - (-1)^{n+1})2^{2n+1}}{(2n+2)!} x^{2n-1} \tan x \\ &< \left(\frac{\sin x}{x} \right)^2 + \frac{\tan x}{x} \\ &< 2 + \sum_{k=3}^n \frac{(2(2^{2k}-1)|B_{2k}| - (-1)^k)2^{2k-1}}{(2k)!} x^{2k-2} + \left(\frac{2}{\pi} \right)^{2n} x^{2n-1} \tan x, \end{aligned}$$

where B_n ($n \in \mathbb{N}_0$, $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$, $\mathbb{N} := \{1, 2, \dots\}$) are the Bernoulli numbers, defined by

$$\frac{t}{e^t - 1} = \sum_{n=0}^{\infty} B_n \frac{t^n}{n!}, \quad |t| < 2\pi.$$

Conjecture 1.2. For $0 < x < \pi/2$ and $n \geq 1$,

$$\left(\frac{x}{\sin x} \right)^2 + \frac{x}{\tan x} < 2 + \sum_{k=2}^n \frac{(k-1) \cdot 2^{2k+1} |B_{2k}|}{(2k)!} x^{2k} + \frac{n \cdot 2^{2n+3} |B_{2(n+1)}|}{(2n+2)!} x^{2n+1} \tan x.$$

Here, and throughout this paper, an empty sum is understood to be zero.

Conjecture 1.3. For $0 < x < \pi/2$ and $n \geq 2$,

$$\begin{aligned}
& 3 + \sum_{k=3}^n \left(\frac{2^{2k}(2^{2k}-1)|B_{2k}|}{4k} - (-1)^k \right) \frac{2}{(2k-1)!} x^{2k-2} \\
& \quad + \left(\frac{2^{2n+2}(2^{2n+2}-1)|B_{2n+2}|}{4(n+1)} - (-1)^{n+1} \right) \frac{2}{(2n+1)!} x^{2n-1} \tan x \\
& < 2 \left(\frac{\sin x}{x} \right) + \frac{\tan x}{x} \\
& < 3 + \sum_{k=3}^n \left(\frac{2^{2k}(2^{2k}-1)|B_{2k}|}{4k} - (-1)^k \right) \frac{2}{(2k-1)!} x^{2k-2} + \left(\frac{2}{\pi} \right)^{2n} x^{2n-1} \tan x.
\end{aligned}$$

Recently, Chen and Paris [4] proved Conjecture 1.2. This paper is a continuation of our earlier work [4]. The first aim of the present paper is to prove Conjectures 1.1 and 1.3.

Mortici [11, Theorem 1] presented the following double inequality:

$$\begin{aligned}
2 + \left(\frac{8}{45} - \frac{8}{945}x^2 \right) x^3 \tan x & < \left(\frac{\sin x}{x} \right)^2 + \frac{\tan x}{x} \\
& < 2 + \left(\frac{8}{45} - \frac{8}{945}x^2 + \frac{16}{14175}x^4 \right) x^3 \tan x, \quad 0 < x < 1.
\end{aligned} \tag{1.11}$$

By using Maple software, we find that

$$\begin{aligned}
\frac{\left(\frac{\sin x}{x} \right)^2 + \frac{\tan x}{x} - 2}{x^3 \tan x} &= \frac{8}{45} - \frac{8}{945}x^2 + \frac{16}{14175}x^4 + \frac{8}{467775}x^6 + \frac{3184}{638512875}x^8 \\
& \quad + \frac{272}{638512875}x^{10} + \frac{7264}{162820783125}x^{12} + \dots.
\end{aligned} \tag{1.12}$$

This fact led us to claim that the upper bound in (1.11) should be the lower bound. The second aim of the present paper is to prove the following inequality:

$$\begin{aligned}
2 + \left(\frac{8}{45} - \frac{8}{945}x^2 + \frac{16}{14175}x^4 \right) x^3 \tan x & < \left(\frac{\sin x}{x} \right)^2 + \frac{\tan x}{x} \\
& < 2 + \left(\frac{8}{45} - \frac{8}{945}x^2 + \frac{241920 - 2688\pi^4 + 32\pi^6}{945\pi^8}x^4 \right) x^3 \tan x, \quad 0 < x < \frac{\pi}{2},
\end{aligned} \tag{1.13}$$

where the constants $\frac{16}{14175}$ and $(241920 - 2688\pi^4 + 32\pi^6)/(945\pi^8)$ are the best possible.

Remark 1.1. The inequalities (1.13) are sharper than the inequalities (1.2) and (1.8).

In analogy with (1.13), we here determine the best possible constants $\alpha, \beta, \lambda, \mu, \rho$, and ϱ such that

$$\begin{aligned}
2 + \left(\frac{2}{45} - \frac{2}{315}x^2 - \alpha x^4 \right) x^3 \tan x & < \left(\frac{x}{\sin x} \right)^2 + \frac{x}{\tan x} < 2 + \left(\frac{2}{45} - \frac{2}{315}x^2 - \beta x^4 \right) x^3 \tan x, \\
3 + \left(\frac{3}{20} + \frac{1}{280}x^2 + \lambda x^4 \right) x^3 \tan x & < 2 \left(\frac{\sin x}{x} \right) + \frac{\tan x}{x} < 3 + \left(\frac{3}{20} + \frac{1}{280}x^2 + \mu x^4 \right) x^3 \tan x
\end{aligned}$$

and

$$3 + \left(\frac{1}{60} - \frac{1}{280}x^2 - \rho x^4 \right) x^3 \tan x < 2 \left(\frac{x}{\sin x} \right) + \frac{x}{\tan x} < 3 + \left(\frac{1}{60} - \frac{1}{280}x^2 - \varrho x^4 \right) x^3 \tan x$$

for $0 < x < \pi/2$. This is the last aim of the present paper.

2. A useful lemma

It is well known that

$$\tan x = \sum_{n=1}^{\infty} \frac{2^{2n}(2^{2n}-1)|B_{2n}|}{(2n)!} x^{2n-1}, \quad |x| < \frac{\pi}{2}, \quad (2.1)$$

By using induction, Chen and Qi [5] (see also [24]) proved the following

Lemma 2.1. *Let $n \geq 1$ be an integer. Then for $0 < x < \pi/2$, we have*

$$\begin{aligned} \frac{2^{2n+2}(2^{2n+2}-1)|B_{2n+2}|}{(2n+2)!} x^{2n} \tan x &< \tan x - \sum_{k=1}^n \frac{2^{2k}(2^{2k}-1)|B_{2k}|}{(2k)!} x^{2k-1} \\ &< \left(\frac{2}{\pi}\right)^{2n} x^{2n} \tan x, \end{aligned} \quad (2.2)$$

where the constants

$$\frac{2^{2n+2}(2^{2n+2}-1)|B_{2n+2}|}{(2n+2)!} \quad \text{and} \quad \left(\frac{2}{\pi}\right)^{2n}$$

are the best possible.

3. Main results

Theorem 3.1. *For $0 < x < \pi/2$ and $n \geq 3$, we have*

$$\begin{aligned} &\left(\frac{(-1)^n 2^{2n+1}}{(2n+2)!} + \frac{2^{2n+2}(2^{2n+2}-1)|B_{2n+2}|}{(2n+2)!} \right) x^{2n-1} \tan x \\ &< \left(\frac{\sin x}{x} \right)^2 + \frac{\tan x}{x} - \left\{ 2 + \sum_{k=3}^n \left(\frac{(-1)^{k-1} 2^{2k-1}}{(2k)!} + \frac{2^{2k}(2^{2k}-1)|B_{2k}|}{(2k)!} \right) x^{2k-2} \right\} \\ &< \left(\frac{2}{\pi} \right)^{2n} x^{2n-1} \tan x. \end{aligned} \quad (3.1)$$

Proof. First of all, we prove the first inequality in (3.1). By using the power series expansions for $\cos x$ and $\tan x$, we have

$$\begin{aligned} \left(\frac{\sin x}{x} \right)^2 + \frac{\tan x}{x} &= \frac{1}{2x^2} - \frac{1}{2x^2} \cos(2x) + \frac{\tan x}{x} \\ &= 2 + \sum_{k=3}^n \left(\frac{(-1)^{k-1} 2^{2k-1}}{(2k)!} + \frac{2^{2k}(2^{2k}-1)|B_{2k}|}{(2k)!} \right) x^{2k-2} + r_n(x), \end{aligned}$$

where

$$r_n(x) = \sum_{k=n+1}^{\infty} \left(\frac{(-1)^{k-1} 2^{2k-1}}{(2k)!} + \frac{2^{2k}(2^{2k}-1)|B_{2k}|}{(2k)!} \right) x^{2k-2}.$$

The first inequality in (3.1) is equivalent to

$$\left(\frac{(-1)^n 2^{2n+1}}{(2n+2)!} + \frac{2^{2n+2}(2^{2n+2}-1)|B_{2n+2}|}{(2n+2)!} \right) x^{2n-1} \tan x < r_n(x)$$

for $0 < x < \pi/2$ and $n \geq 3$, which can be written by (2.1) as

$$\sum_{k=n+2}^{\infty} \left\{ \left(\frac{(-1)^n 2^{2n+1}}{(2n+2)!} + \frac{2^{2n+2}(2^{2n+2}-1)|B_{2n+2}|}{(2n+2)!} \right) \frac{2^{2k-2n}(2^{2k-2n}-1)|B_{2k-2n}|}{(2k-2n)!} \right. \\ \left. - \left(\frac{(-1)^{k-1} 2^{2k-1}}{(2k)!} + \frac{2^{2k}(2^{2k}-1)|B_{2k}|}{(2k)!} \right) \right\} x^{2k-2} < 0,$$

where we note that the term corresponding to $k = n+1$ vanishes.

We claim that for $k \geq n+2$,

$$\left(\frac{(-1)^n 2^{2n+1}}{(2n+2)!} + \frac{2^{2n+2}(2^{2n+2}-1)|B_{2n+2}|}{(2n+2)!} \right) \frac{2^{2k-2n}(2^{2k-2n}-1)|B_{2k-2n}|}{(2k-2n)!} \\ < \frac{(-1)^{k-1} 2^{2k-1}}{(2k)!} + \frac{2^{2k}(2^{2k}-1)|B_{2k}|}{(2k)!}, \quad (3.2)$$

It is enough to prove the following inequality:

$$\left(\frac{2^{2n+1}}{(2n+2)!} + \frac{2^{2n+2}(2^{2n+2}-1)|B_{2n+2}|}{(2n+2)!} \right) \frac{2^{2k-2n}(2^{2k-2n}-1)|B_{2k-2n}|}{(2k-2n)!} \\ < -\frac{2^{2k-1}}{(2k)!} + \frac{2^{2k}(2^{2k}-1)|B_{2k}|}{(2k)!}, \quad k \geq n+2.$$

Using the following inequality (see [7]):

$$\frac{2}{(2\pi)^{2n}(1-2^{-2n})} < \frac{|B_{2n}|}{(2n)!} < \frac{2}{(2\pi)^{2n}(1-2^{1-2n})}, \quad n \geq 1, \quad (3.3)$$

it suffices to show that for $k \geq n+2$,

$$\left(\frac{2^{2n+1}}{(2n+2)!} + \frac{2^{2n+2}(2^{2n+2}-1)2}{(2\pi)^{2n+2}(1-2^{1-2(n+1)})} \right) \frac{2^{2k-2n}(2^{2k-2n}-1)2}{(2\pi)^{2k-2n}(1-2^{1-2(k-n)})} \\ < -\frac{2^{2k-1}}{(2k)!} + \frac{2^{2k}(2^{2k}-1)2}{(2\pi)^{2k}(1-2^{-2k})},$$

which can be rearranged as

$$\left(\frac{2^{2n+1}}{(2n+2)!} + \frac{2(2^{2n+2}-1)}{2^{2n+2}-2} \left(\frac{2}{\pi} \right)^{2n+2} \right) \frac{2^{2k-2n}-1}{2^{2k-2n}-2} \left(\frac{2}{\pi} \right)^{2k-2n} + \frac{2^{2k-2}}{(2k)!} < \left(\frac{2}{\pi} \right)^{2k}, \\ \left(\frac{2^{2n+1}}{(2n+2)!} + 2 \left(1 + \frac{1}{2^{2n+2}-2} \right) \left(\frac{2}{\pi} \right)^{2n+2} \right) \left(1 + \frac{1}{2^{2k-2n}-2} \right) \left(\frac{\pi}{2} \right)^{2n} \\ + \frac{2^{2k-2}}{(2k)!} \left(\frac{\pi}{2} \right)^{2k} < 1, \quad k \geq n+2.$$

Noting that the sequences

$$1 + \frac{1}{2^{2k-2n}-2} \quad \text{and} \quad \frac{2^{2k-2}}{(2k)!} \left(\frac{\pi}{2} \right)^{2k}$$

are both strictly decreasing for $k \geq n+2$, it suffices to show that

$$\left(\frac{2^{2n+1}}{(2n+2)!} + 2 \left(1 + \frac{1}{2^{2n+2}-2} \right) \left(\frac{2}{\pi} \right)^{2n+2} \right) \frac{15}{14} \left(\frac{\pi}{2} \right)^{2n} + \frac{2^{2n+2}}{(2n+4)!} \left(\frac{\pi}{2} \right)^{2n+4} < 1$$

for $n \geq 3$, which can be rearranged as

$$\frac{2^{2n+1}}{(2n+2)!} \left(\frac{\pi}{2}\right)^{2n+2} + \frac{1}{2^{2n+1}-1} + \frac{2^{2n+2}}{(2n+4)!} \left(\frac{\pi}{2}\right)^{2n+6} \left(\frac{14}{15}\right) < \frac{14}{15} \left(\frac{\pi}{2}\right)^2 - 2, \quad n \geq 3.$$

Noting that the sequence

$$a_n := \frac{2^{2n+1}}{(2n+2)!} \left(\frac{\pi}{2}\right)^{2n+2} + \frac{1}{2^{2n+1}-1} + \frac{2^{2n+2}}{(2n+4)!} \left(\frac{\pi}{2}\right)^{2n+6} \left(\frac{14}{15}\right), \quad n \geq 3$$

is strictly decreasing, we see that

$$a_n \leq a_3 = \frac{1}{127} + \frac{\pi^8}{80640} + \frac{\pi^{12}}{62208000} < \frac{14}{15} \left(\frac{\pi}{2}\right)^2 - 2$$

holds true for $n \geq 3$, since

$$\frac{1}{127} + \frac{\pi^8}{80640} + \frac{\pi^{12}}{62208000} = 0.14039705\dots, \quad \frac{14}{15} \left(\frac{\pi}{2}\right)^2 - 2 = 0.30290769\dots$$

This proves the claim (3.2). Hence, the first inequality in (3.1) holds for $0 < x < \pi/2$ and $n \geq 3$.

Secondly, we prove the second inequality in (3.1). We consider two cases.

Case 1. $n = 2N + 1$ ($N \geq 1$).

It is well known that for $x \neq 0$,

$$\sum_{k=1}^{2N} (-1)^{k-1} \frac{x^{2k-2}}{(2k-2)!} < \cos x < \sum_{k=1}^{2N+1} (-1)^{k-1} \frac{x^{2k-2}}{(2k-2)!}.$$

We then obtain that

$$\left(\frac{\sin x}{x}\right)^2 = \frac{1}{2x^2} - \frac{1}{2x^2} \cos(2x) < \sum_{k=1}^{2N+1} (-1)^{k-1} \frac{2^{2k-1}}{(2k)!} x^{2k-2}. \quad (3.4)$$

The choice $n = 2N + 1$ in (2.2), we obtain from the right-hand inequality of (2.2) that

$$\frac{\tan x}{x} < \sum_{k=1}^{2N+1} \frac{2^{2k}(2^{2k}-1)|B_{2k}|}{(2k)!} x^{2k-2} + \left(\frac{2}{\pi}\right)^{4N+2} x^{4N+1} \tan x. \quad (3.5)$$

Adding these two expressions, we obtain

$$\begin{aligned} \left(\frac{\sin x}{x}\right)^2 + \frac{\tan x}{x} &< 2 + \sum_{k=3}^{2N+1} \left(\frac{(-1)^{k-1} 2^{2k-1}}{(2k)!} + \frac{2^{2k}(2^{2k}-1)|B_{2k}|}{(2k)!} \right) x^{2k-2} \\ &\quad + \left(\frac{2}{\pi}\right)^{4N+2} x^{4N+1} \tan x. \end{aligned}$$

This shows that the second inequality in (3.1) holds for $n = 2N + 1$.

Case 2. $n = 2N$ ($N \geq 2$).

Write

$$\begin{aligned} \left(\frac{\sin x}{x}\right)^2 + \frac{\tan x}{x} &= 2 + \sum_{k=3}^{2N} \left(\frac{(-1)^{k-1} 2^{2k-1}}{(2k)!} + \frac{2^{2k}(2^{2k}-1)|B_{2k}|}{(2k)!} \right) x^{2k-2} \\ &\quad + \sum_{k=2N+1}^{\infty} \left(\frac{(-1)^{k-1} 2^{2k-1}}{(2k)!} + \frac{2^{2k}(2^{2k}-1)|B_{2k}|}{(2k)!} \right) x^{2k-2}. \end{aligned}$$

We need to prove

$$\sum_{k=2N+1}^{\infty} \left(\frac{(-1)^{k-1} 2^{2k-1}}{(2k)!} + \frac{2^{2k}(2^{2k}-1)|B_{2k}|}{(2k)!} \right) x^{2k-2} < \left(\frac{2}{\pi} \right)^{4N} x^{4N-1} \tan x. \quad (3.6)$$

Noting that (2.1) holds, we can rewrite (3.6) as

$$\sum_{k=2N+1}^{\infty} \left\{ \frac{(-1)^{k-1} 2^{2k-1}}{(2k)!} + \frac{2^{2k}(2^{2k}-1)|B_{2k}|}{(2k)!} - \left(\frac{2}{\pi} \right)^{4N} \frac{2^{2k-4N}(2^{2k-4N}-1)|B_{2k-4N}|}{(2k-4N)!} \right\} x^{2k-2} < 0.$$

We claim that for $k \geq 2N+1$,

$$\frac{(-1)^{k-1} 2^{2k-1}}{(2k)!} + \frac{2^{2k}(2^{2k}-1)|B_{2k}|}{(2k)!} < \left(\frac{2}{\pi} \right)^{4N} \frac{2^{2k-4N}(2^{2k-4N}-1)|B_{2k-4N}|}{(2k-4N)!}. \quad (3.7)$$

It is enough to prove the following inequality:

$$\frac{2^{2k-1}}{(2k)!} + \frac{2^{2k}(2^{2k}-1)|B_{2k}|}{(2k)!} < \left(\frac{2}{\pi} \right)^{4N} \frac{2^{2k-4N}(2^{2k-4N}-1)|B_{2k-4N}|}{(2k-4N)!}, \quad k \geq 2N+1. \quad (3.8)$$

Using (3.3), we find that for $k \geq 2N+1$,

$$\frac{\frac{2^{2k+2}(2^{2k+2}-1)|B_{2k+2}|}{(2k+2)!}}{\frac{2^{2k}(2^{2k}-1)|B_{2k}|}{(2k)!}} < \frac{\frac{2^{2k+2}(2^{2k+2}-1)2}{(2\pi)^{2k+2}(1-2^{1-2(k+1)})}}{\frac{2^{2k}(2^{2k}-1)2}{(2\pi)^{2k}(1-2^{-2k})}} = \frac{2(4^k-2)(4 \cdot 4^k-1)}{\pi^2(4^k-1)(2 \cdot 4^k-1)} < 1 \quad (3.9)$$

and¹

$$\begin{aligned} & \frac{\frac{2^{2k-4N+2}(2^{2k-4N+2}-1)|B_{2k-4N+2}|}{(2k-4N+2)!}}{\frac{2^{2k-4N}(2^{2k-4N}-1)|B_{2k-4N}|}{(2k-4N)!}} > \frac{\frac{2^{2k-4N+2}(2^{2k-4N+2}-1)2}{(2\pi)^{2k-4N+2}(1-2^{-2(k-2N+1)})}}{\frac{2^{2k-4N}(2^{2k-4N}-1)2}{(2\pi)^{2k-4N}(1-2^{1-2(k-2N)})}} \\ & = \frac{16^{k+N+1} - (8 \cdot 256^N + 64^N)4^{k+1} + 8 \cdot 1024^N}{\pi^2(4^k-16^N)(4^{k+1}-16^N)} > 1. \end{aligned} \quad (3.10)$$

Hence, the sequence

$$\frac{2^{2k-1}}{(2k)!} + \frac{2^{2k}(2^{2k}-1)|B_{2k}|}{(2k)!}$$

is strictly decreasing, and the sequence

$$\left(\frac{2}{\pi} \right)^{2N} \frac{2^{2k-2N}(2^{2k-2N}-1)|B_{2k-2N}|}{(2k-2N)!}$$

¹The inequality (3.10) is proved in the appendix.

is strictly increasing for $k \geq 2N+1$. In order to prove (3.8), it suffices to show that for $k \geq 2N+1$,

$$\frac{2^{4N+1}}{(4N+2)!} + \frac{2^{4N+2}(2^{4N+2}-1)|B_{4N+2}|}{(4N+2)!} < \left(\frac{2}{\pi}\right)^{4N} \frac{2^2(2^2-1)|B_2|}{2!} = \left(\frac{2}{\pi}\right)^{4N}. \quad (3.11)$$

By (3.3), it suffices to show that

$$\begin{aligned} \frac{2^{4N+1}}{(4N+2)!} + \frac{2^{4N+2}(2^{4N+2}-1)2}{(2\pi)^{4N+2}(1-2^{1-2(2N+1)})} &< \left(\frac{2}{\pi}\right)^{4N}, \\ \frac{2^{4N+1}}{(4N+2)!} + 2\left(1 + \frac{1}{2^{4N+2}-2}\right) \left(\frac{2}{\pi}\right)^{4N+2} &< \left(\frac{2}{\pi}\right)^{4N}, \\ \frac{2^{4N+1}}{(4N+2)!} \left(\frac{\pi}{2}\right)^{4N+2} + \frac{1}{2^{4N+1}-1} &< \left(\frac{\pi}{2}\right)^2 - 2, \quad N \geq 2. \end{aligned}$$

Noting that the sequence

$$b_N := \frac{2^{4N+1}}{(4N+2)!} \left(\frac{\pi}{2}\right)^{4N+2} + \frac{1}{2^{4N+1}-1}, \quad N \geq 2$$

is strictly decreasing, we see that

$$b_N \leq b_2 = \frac{1}{511} + \frac{\pi^{10}}{7257600} < \left(\frac{\pi}{2}\right)^2 - 2$$

holds true for $N \geq 2$, since

$$\frac{1}{511} + \frac{\pi^{10}}{7257600} = 0.0148603\dots, \quad \left(\frac{\pi}{2}\right)^2 - 2 = 0.4674011\dots$$

This proves the claim (3.7). Hence, (3.6) holds, which shows that the second inequality in (3.1) holds for $n = 2N$. Thus, the second inequality in (3.1) holds for $0 < x < \pi/2$ and $n \geq 3$. The proof of Theorem 3.1 is complete. $\square$

Theorem 3.2. For $0 < x < \pi/2$ and $n \geq 2$, we have

$$\begin{aligned} &\left(\frac{2(-1)^n}{(2n+1)!} + \frac{2^{2n+2}(2^{2n+2}-1)|B_{2n+2}|}{(2n+2)!} \right) x^{2n-1} \tan x \\ &< 2 \left(\frac{\sin x}{x} \right) + \frac{\tan x}{x} - \left\{ 3 + \sum_{k=3}^n \left(\frac{2(-1)^{k-1}}{(2k-1)!} + \frac{2^{2k}(2^{2k}-1)|B_{2k}|}{(2k)!} \right) x^{2k-2} \right\} \\ &< \left(\frac{2}{\pi} \right)^{2n} x^{2n-1} \tan x. \end{aligned} \quad (3.12)$$

Proof. First of all, we prove the first inequality in (3.12). By using the power series expansions for $\sin x$ and $\tan x$, we have

$$2 \left(\frac{\sin x}{x} \right) + \frac{\tan x}{x} = 3 + \sum_{k=3}^n \left(\frac{2(-1)^{k-1}}{(2k-1)!} + \frac{2^{2k}(2^{2k}-1)|B_{2k}|}{(2k)!} \right) x^{2k-2} + R_n(x),$$

where

$$R_n(x) = \sum_{k=n+1}^{\infty} \left(\frac{2(-1)^{k-1}}{(2k-1)!} + \frac{2^{2k}(2^{2k}-1)|B_{2k}|}{(2k)!} \right) x^{2k-2}.$$

The first inequality in (3.12) is equivalent to

$$\left(\frac{2(-1)^n}{(2n+1)!} + \frac{2^{2n+2}(2^{2n+2}-1)|B_{2n+2}|}{(2n+2)!} \right) x^{2n-1} \tan x < R_n(x)$$

for $0 < x < \pi/2$ and $n \geq 2$, which can be written by (2.1) as

$$\sum_{k=n+2}^{\infty} \left\{ \left(\frac{2(-1)^n}{(2n+1)!} + \frac{2^{2n+2}(2^{2n+2}-1)|B_{2n+2}|}{(2n+2)!} \right) \frac{2^{2k-2n}(2^{2k-2n}-1)|B_{2k-2n}|}{(2k-2n)!} \right. \\ \left. - \left(\frac{2(-1)^{k-1}}{(2k-1)!} + \frac{2^{2k}(2^{2k}-1)|B_{2k}|}{(2k)!} \right) \right\} x^{2k-2} < 0,$$

where we note that the term corresponding to $k = n+1$ vanishes.

We claim that for $k \geq n+2$,

$$\left(\frac{2(-1)^n}{(2n+1)!} + \frac{2^{2n+2}(2^{2n+2}-1)|B_{2n+2}|}{(2n+2)!} \right) \frac{2^{2k-2n}(2^{2k-2n}-1)|B_{2k-2n}|}{(2k-2n)!} \\ < \frac{2(-1)^{k-1}}{(2k-1)!} + \frac{2^{2k}(2^{2k}-1)|B_{2k}|}{(2k)!}. \quad (3.13)$$

It is enough to prove the following inequality:

$$\left(\frac{2}{(2n+1)!} + \frac{2^{2n+2}(2^{2n+2}-1)|B_{2n+2}|}{(2n+2)!} \right) \frac{2^{2k-2n}(2^{2k-2n}-1)|B_{2k-2n}|}{(2k-2n)!} \\ < -\frac{2}{(2k-1)!} + \frac{2^{2k}(2^{2k}-1)|B_{2k}|}{(2k)!}, \quad k \geq n+2.$$

Using (3.3), it suffices to show that for $k \geq n+2$,

$$\left(\frac{2}{(2n+1)!} + \frac{2^{2n+2}(2^{2n+2}-1)2}{(2\pi)^{2n+2}(1-2^{1-2(n+1)})} \right) \frac{2^{2k-2n}(2^{2k-2n}-1)2}{(2\pi)^{2k-2n}(1-2^{1-2(k-n)})} \\ < -\frac{2}{(2k-1)!} + \frac{2^{2k}(2^{2k}-1)2}{(2\pi)^{2k}(1-2^{-2k})},$$

which can be rearranged as

$$\left(\frac{1}{(2n+1)!} + \frac{2^{2n+2}-1}{2^{2n+2}-2} \left(\frac{2}{\pi} \right)^{2n+2} \right) \frac{2(2^{2k-2n}-1)}{2^{2k-2n}-2} \left(\frac{2}{\pi} \right)^{2k-2n} + \frac{1}{(2k-1)!} < \left(\frac{2}{\pi} \right)^{2k}, \\ \left(\frac{1}{(2n+1)!} + \left(1 + \frac{1}{2^{2n+2}-2} \right) \left(\frac{2}{\pi} \right)^{2n+2} \right) 2 \left(1 + \frac{1}{2^{2k-2n}-2} \right) \left(\frac{\pi}{2} \right)^{2n} \\ + \frac{1}{(2k-1)!} \left(\frac{\pi}{2} \right)^{2k} < 1, \quad k \geq n+2.$$

Noting that the sequences

$$2 \left(1 + \frac{1}{2^{2k-2n}-2} \right) \quad \text{and} \quad \frac{1}{(2k-1)!} \left(\frac{\pi}{2} \right)^{2k}$$

are both strictly decreasing for $k \geq n+2$, it suffices to show that

$$\left(\frac{1}{(2n+1)!} + \left(1 + \frac{1}{2^{2n+2}-2} \right) \left(\frac{2}{\pi} \right)^{2n+2} \right) \frac{15}{7} \left(\frac{\pi}{2} \right)^{2n} + \frac{1}{(2n+3)!} \left(\frac{\pi}{2} \right)^{2n+4} < 1$$

for $n \geq 2$, which can be rearranged as

$$\frac{1}{(2n+1)!} \left(\frac{\pi}{2}\right)^{2n+2} + \frac{1}{2^{2n+2}-2} + \frac{1}{(2n+3)!} \left(\frac{\pi}{2}\right)^{2n+6} \left(\frac{7}{15}\right) < \frac{7}{15} \left(\frac{\pi}{2}\right)^2 - 1, \quad n \geq 2.$$

Noting that the sequence

$$x_n := \frac{1}{(2n+1)!} \left(\frac{\pi}{2}\right)^{2n+2} + \frac{1}{2^{2n+2}-2} + \frac{1}{(2n+3)!} \left(\frac{\pi}{2}\right)^{2n+6} \left(\frac{7}{15}\right), \quad n \geq 2$$

is strictly decreasing, we see that

$$x_n \leq x_2 = \frac{1}{62} + \frac{\pi^6}{7680} + \frac{\pi^{10}}{11059200} < \frac{7}{15} \left(\frac{\pi}{2}\right)^2 - 1$$

holds true for $n \geq 2$, since

$$\frac{1}{62} + \frac{\pi^6}{7680} + \frac{\pi^{10}}{11059200} = 0.1497778\dots, \quad \frac{7}{15} \left(\frac{\pi}{2}\right)^2 - 1 = 0.1514538\dots$$

This proves the claim (3.13). Hence, the first inequality in (3.12) holds for $0 < x < \pi/2$ and $n \geq 2$.

Secondly, we prove the second inequality in (3.12). We consider two cases.

Case 1. $n = 2N + 1$ ($N \geq 1$).

It is well known that for $x \neq 0$,

$$\sum_{k=1}^{2N} (-1)^{k-1} \frac{x^{2k-2}}{(2k-1)!} < \frac{\sin x}{x} < \sum_{k=1}^{2N+1} (-1)^{k-1} \frac{x^{2k-2}}{(2k-1)!}. \quad (3.14)$$

From the second inequality in (3.14) and (3.5), we obtain

$$\begin{aligned} 2 \left(\frac{\sin x}{x} \right) + \frac{\tan x}{x} &< 3 + \sum_{k=3}^{2N+1} \left(\frac{2(-1)^{k-1}}{(2k-1)!} + \frac{2^{2k}(2^{2k}-1)|B_{2k}|}{(2k)!} \right) x^{2k-2} \\ &\quad + \left(\frac{2}{\pi} \right)^{4N+2} x^{4N+1} \tan x. \end{aligned}$$

This shows that the second inequality in (3.12) holds for $n = 2N + 1$.

Case 2. $n = 2N$ ($N \geq 1$).

Write

$$\begin{aligned} 2 \left(\frac{\sin x}{x} \right) + \frac{\tan x}{x} &= 3 + \sum_{k=3}^{2N} \left(\frac{2(-1)^{k-1}}{(2k-1)!} + \frac{2^{2k}(2^{2k}-1)|B_{2k}|}{(2k)!} \right) x^{2k-2} \\ &\quad + \sum_{k=2N+1}^{\infty} \left(\frac{2(-1)^{k-1}}{(2k-1)!} + \frac{2^{2k}(2^{2k}-1)|B_{2k}|}{(2k)!} \right) x^{2k-2}. \end{aligned}$$

We need to prove

$$\sum_{k=2N+1}^{\infty} \left(\frac{2(-1)^{k-1}}{(2k-1)!} + \frac{2^{2k}(2^{2k}-1)|B_{2k}|}{(2k)!} \right) x^{2k-2} < \left(\frac{2}{\pi} \right)^{4N} x^{4N-1} \tan x. \quad (3.15)$$

Noting that (2.1) holds, we can rewrite (3.15) as

$$\sum_{k=2N+1}^{\infty} \left\{ \frac{2(-1)^{k-1}}{(2k-1)!} + \frac{2^{2k}(2^{2k}-1)|B_{2k}|}{(2k)!} - \left(\frac{2}{\pi}\right)^{4N} \frac{2^{2k-4N}(2^{2k-4N}-1)|B_{2k-4N}|}{(2k-4N)!} \right\} x^{2k-2} < 0.$$

We claim that for $k \geq 2N+1$,

$$\frac{2(-1)^{k-1}}{(2k-1)!} + \frac{2^{2k}(2^{2k}-1)|B_{2k}|}{(2k)!} < \left(\frac{2}{\pi}\right)^{4N} \frac{2^{2k-4N}(2^{2k-4N}-1)|B_{2k-4N}|}{(2k-4N)!}. \quad (3.16)$$

It is enough to prove the following inequality:

$$\frac{2}{(2k-1)!} + \frac{2^{2k}(2^{2k}-1)|B_{2k}|}{(2k)!} < \left(\frac{2}{\pi}\right)^{4N} \frac{2^{2k-4N}(2^{2k-4N}-1)|B_{2k-4N}|}{(2k-4N)!}, \quad k \geq 2N+1. \quad (3.17)$$

By (3.9) and (3.10), we see that the sequence

$$\frac{2}{(2k-1)!} + \frac{2^{2k}(2^{2k}-1)|B_{2k}|}{(2k)!}$$

is strictly decreasing, and the sequence

$$\left(\frac{2}{\pi}\right)^{2N} \frac{2^{2k-2N}(2^{2k-2N}-1)|B_{2k-2N}|}{(2k-2N)!}$$

is strictly increasing for $k \geq 2N+1$. In order to prove (3.17), it suffices to show that for $k \geq 2N+1$,

$$\frac{2}{(4N+1)!} + \frac{2^{4N+2}(2^{4N+2}-1)|B_{4N+2}|}{(4N+2)!} < \left(\frac{2}{\pi}\right)^{4N} \frac{2^2(2^2-1)|B_2|}{2!} = \left(\frac{2}{\pi}\right)^{4N}. \quad (3.18)$$

By (3.3), it now suffices to show that

$$\frac{2}{(4N+1)!} + \frac{2^{4N+2}(2^{4N+2}-1)2}{(2\pi)^{4N+2}(1-2^{1-2(2N+1)})} < \left(\frac{2}{\pi}\right)^{4N},$$

which can be rearranged as

$$\left(\frac{2}{\pi}\right)^{4N+2} \frac{1}{(4N+1)!} + \frac{1}{2^{4N+2}-2} < \frac{\pi^2}{8} - 1, \quad N \geq 1.$$

Noting that the sequence

$$y_N := \left(\frac{2}{\pi}\right)^{4N+2} \frac{1}{(4N+1)!} + \frac{1}{2^{4N+2}-2}, \quad N \geq 1$$

is strictly decreasing, we see that

$$y_N \leq y_1 = \frac{\pi^4}{48384} < \frac{\pi^2}{8} - 1$$

holds true for $N \geq 1$, since

$$\frac{\pi^4}{48384} = 0.00201325\dots, \quad \frac{\pi^2}{8} - 1 = 0.23370055\dots$$

This proves the claim (3.16). Hence, (3.15) holds, which shows that the second inequality in (3.12) holds for $n = 2N$. Thus, the second inequality in (3.12) holds for $0 < x < \pi/2$ and $n \geq 2$. The proof of Theorem 3.2 is complete. $\square$

Theorem 3.3. *For $0 < x < \pi/2$, we have*

$$\begin{aligned} 2 + \left(\frac{8}{45} - \frac{8}{945}x^2 + ax^4 \right) x^3 \tan x &< \left(\frac{\sin x}{x} \right)^2 + \frac{\tan x}{x} \\ &< 2 + \left(\frac{8}{45} - \frac{8}{945}x^2 + bx^4 \right) x^3 \tan x \end{aligned} \quad (3.19)$$

with the best possible constants

$$a = \frac{16}{14175} = 0.001128\dots \quad \text{and} \quad b = \frac{241920 - 2688\pi^4 + 32\pi^6}{945\pi^8} = 0.001209\dots \quad (3.20)$$

Proof. The inequality (3.19) can be written as

$$a < f(x) < b,$$

where

$$f(x) = \frac{1}{x^4} \left(\frac{\left(\frac{\sin x}{x} \right)^2 + \frac{\tan x}{x} - 2}{x^3 \tan x} - \left(\frac{8}{45} - \frac{8}{945}x^2 \right) \right).$$

Direct computations yield

$$\lim_{x \rightarrow 0} f(x) = \frac{16}{14175} \quad \text{and} \quad \lim_{x \rightarrow \pi/2} f(x) = \frac{241920 - 2688\pi^4 + 32\pi^6}{945\pi^8}.$$

In order to prove (3.19), it suffices to show that $f(x)$ is strictly increasing on $(0, \pi/2)$.

Differentiation yields

$$f'(x) = \frac{g(x)}{945x^{10} \sin^2 x},$$

with

$$\begin{aligned} g(x) &= 6615x^2 \sin(2x) - 8505 \sin^3 x \cos x - 8505x + 1890x^3 \\ &\quad + 10395x \cos^2 x - 1890x \cos^4 x + 672x^5 \sin^2 x - 16x^7 \sin^2 x \\ &= 6615x^2 \sin(2x) - 8505 \left(\frac{1}{4} \sin(2x) - \frac{1}{8} \sin(4x) \right) - 8505x + 1890x^3 \\ &\quad + \frac{10395}{2}x \left(1 + \cos(2x) \right) - 1890x \left(\frac{1}{8} \cos(4x) + \frac{1}{2} \cos(2x) + \frac{3}{8} \right) \\ &\quad + 336x^5 \left(1 - \cos(2x) \right) - 8x^7 \left(1 - \cos(2x) \right) \\ &= \frac{16}{495}x^{13} + \frac{496}{61425}x^{15} - \frac{64}{26325}x^{17} + \sum_{n=9}^{\infty} (-1)^{n-1} u_n(x), \end{aligned}$$

where

$$\begin{aligned} u_n(x) &= \left(945n \cdot 2^{2n-1} - 16065 \cdot 2^{2n-2} + 16n^7 - 112n^6 + 952n^5 \right. \\ &\quad \left. - 1960n^4 + 889n^3 + 13727n^2 - 2172n \right) \frac{2^{2n} x^{2n+1}}{(2n+1)!}, \quad n \geq 9. \end{aligned}$$

Direct computation yields

$$\frac{u_{n+1}(x)}{u_n(x)} = \frac{8x^2 p_n}{q_n},$$

where

$$p_n = (1890n - 16065)4^n + 16n^7 + 616n^5 + 1680n^4 + 889n^3 + 12810n^2 + 24309n + 11340$$

and

$$q_n = (n+1)(2n+3) \left((1890n - 16065)4^n + 64n^7 - 448n^6 + 3808n^5 - 7840n^4 + 3556n^3 + 54908n^2 - 8688n \right).$$

Noting that $8(\pi/2)^2 < 20$, we find that for $0 < x < \pi/2$ and $n \geq 9$,

$$\frac{u_{n+1}(x)}{u_n(x)} < \frac{8(\pi/2)^2 p_n}{q_n} < \frac{20p_n}{q_n} < 1,$$

since²

$$q_n - 20p_n > 0 \quad \text{for } n \geq 9. \quad (3.21)$$

Therefore, for fixed $x \in (0, \pi/2)$, the sequence $n \mapsto u_n(x)$ is strictly decreasing for $n \geq 9$. Hence, for $0 < x < \pi/2$,

$$g(x) > \frac{16}{495}x^{13} + \frac{496}{61425}x^{15} - \frac{64}{26325}x^{17} = \frac{16}{495}x^{13} + \frac{16x^{15}(93 - 28x^2)}{184275} > 0.$$

We then obtain that $f'(x) > 0$ for $0 < x < \pi/2$. The proof of Theorem 3.3 is complete. $\square$

Following the same method used in the proof of Theorem 3.3, we can prove the following theorem.

Theorem 3.4. *For $0 < x < \pi/2$, we have*

$$\begin{aligned} 2 + \left(\frac{2}{45} - \frac{2}{315}x^2 - \alpha x^4 \right) x^3 \tan x &< \left(\frac{x}{\sin x} \right)^2 + \frac{x}{\tan x} \\ &< 2 + \left(\frac{2}{45} - \frac{2}{315}x^2 - \beta x^4 \right) x^3 \tan x, \end{aligned} \quad (3.22)$$

$$\begin{aligned} 3 + \left(\frac{3}{20} + \frac{1}{280}x^2 + \lambda x^4 \right) x^3 \tan x &< 2 \left(\frac{\sin x}{x} \right) + \frac{\tan x}{x} \\ &< 3 + \left(\frac{3}{20} + \frac{1}{280}x^2 + \mu x^4 \right) x^3 \tan x \end{aligned} \quad (3.23)$$

and

$$\begin{aligned} 3 + \left(\frac{1}{60} - \frac{1}{280}x^2 - \rho x^4 \right) x^3 \tan x &< 2 \left(\frac{x}{\sin x} \right) + \frac{x}{\tan x} \\ &< 3 + \left(\frac{1}{60} - \frac{1}{280}x^2 - \varrho x^4 \right) x^3 \tan x, \end{aligned} \quad (3.24)$$

with the best possible constants

$$\alpha = \frac{224 - 8\pi^2}{315\pi^4} = 0.004727\dots, \quad \beta = \frac{4}{1575} = 0.002539\dots, \quad (3.25)$$

²The inequality (3.21) is proved in the appendix.

$$\lambda = \frac{23}{33600} = 0.000684\dots, \quad \mu = \frac{17920 - 168\pi^4 - \pi^6}{70\pi^8} = 0.000894\dots \quad (3.26)$$

and

$$\rho = \frac{56 - 3\pi^2}{210\pi^4} = 0.0012901\dots, \quad \varrho = \frac{83}{100800} = 0.0008234\dots \quad (3.27)$$

Proof. We only prove inequality (3.24). The proofs of (3.22) and (3.23) are analogous. The inequality (3.24) can be written as

$$\rho > F(x) > \varrho,$$

where

$$F(x) = \frac{1}{x^4} \left(-\frac{2\left(\frac{x}{\sin x}\right) + \frac{x}{\tan x} - 3}{x^3 \tan x} + \left(\frac{1}{60} - \frac{1}{280}x^2\right) \right).$$

Direct computations yield

$$\lim_{x \rightarrow 0} F(x) = \frac{83}{100800} \quad \text{and} \quad \lim_{x \rightarrow \pi/2} F(x) = \frac{56 - 3\pi^2}{210\pi^4}.$$

In order to prove (3.24), it suffices to show that $F(x)$ is strictly increasing on $(0, \pi/2)$.

Differentiation yields

$$F'(x) = \frac{G(x)}{420x^8 \sin^3 x},$$

with

$$\begin{aligned} G(x) &= 2520x \sin(2x) + (2520x - 3x^5 + 28x^3) \sin x \cos^2 x \\ &\quad + (3x^5 - 1260x - 28x^3) \sin x + (840x^2 - 8820) \cos x + 840x^2 \cos^2 x \\ &\quad + 8820 \cos^3 x + 840x^2 \\ &= 2520x \sin(2x) + (2520x - 3x^5 + 28x^3) \left(\frac{1}{4} \sin x + \frac{1}{4} \sin(3x) \right) \\ &\quad + (3x^5 - 1260x - 28x^3) \sin x + (840x^2 - 8820) \cos x + 420x^2 (1 + \cos(2x)) \\ &\quad + 8820 \left(\frac{1}{4} \cos(3x) + \frac{3}{4} \cos x \right) + 840x^2 \\ &= \sum_{n=6}^{\infty} (-1)^n U_n(x), \end{aligned}$$

where

$$\begin{aligned} U_n(x) &= \left((8n^5 - 40n^4 + 238n^3 - 302n^2 - 33924n + 178605)9^n - (34020n^2 + 187110n)4^n \right. \\ &\quad \left. - 5832n^5 + 29160n^4 - 64638n^3 - 215298n^2 + 222588n - 178605 \right) \frac{x^{2n}}{81 \cdot (2n)!}. \end{aligned}$$

Direct computation yields

$$\frac{U_{n+1}(x)}{U_n(x)} = \frac{9x^2 P_n}{2Q_n},$$

where

$$P_n = (8n^5 + 158n^3 + 252n^2 - 33934n + 144585)9^n - (15120n^2 + 113400n + 98280)4^n - 648n^5 - 32508n^2 - 34938n - 23625 - 702n^3$$

and

$$Q_n = (2n+1)(n+1) \left((8n^5 - 40n^4 + 238n^3 - 302n^2 - 33924n + 178605)9^n - (34020n^2 + 187110n)4^n - 5832n^5 + 29160n^4 - 64638n^3 - 215298n^2 + 222588n - 178605 \right).$$

Noting that $\frac{9}{2} \left(\frac{\pi}{2}\right)^2 < 12$, we find that for $0 < x < \pi/2$ and $n \geq 6$,

$$\frac{U_{n+1}(x)}{U_n(x)} < \frac{9(\pi/2)^2 P_n}{2Q_n} < \frac{12P_n}{Q_n} < 1,$$

since³

$$Q_n - 12P_n > 0 \quad \text{for } n \geq 6. \quad (3.28)$$

Therefore, for fixed $x \in (0, \pi/2)$, the sequence $n \mapsto U_n(x)$ is strictly decreasing for $n \geq 6$. Hence, we have

$$G(x) > 0, \quad 0 < x < \frac{\pi}{2}.$$

We then obtain that $F'(x) > 0$ for $0 < x < \pi/2$. Hence, the inequality (3.24) holds with the best possible constants given in (3.27). The proof is complete. $\square$

Remark 3.1. *The upper bound in (3.22) is sharper than the upper bound in (1.9). The inequalities (3.23) are sharper than the inequalities (1.10).*

Remark 3.2. *Chen and Paris [4] showed that for $0 < x < \pi/2$,*

$$3 + \theta_1 x^3 \tan x < 2 \left(\frac{x}{\sin x} \right) + \frac{x}{\tan x} < 3 + \theta_2 x^3 \tan x \quad (3.29)$$

with the best possible constants

$$\theta_1 = 0 \quad \text{and} \quad \theta_2 = \frac{1}{60}.$$

The double inequality (3.24) is an improvement on the double inequality (3.29).

Appendix A: Proof of (3.10)

Noting that $\pi^2 < 10$, in order to prove (3.10), it suffices to show that for $k \geq 2N + 1$,

$$\begin{aligned} & 16^{k+N+1} - (8 \cdot 256^N + 64^N)4^{k+1} + 8 \cdot 1024^N - 10(4^k - 16^N)(4^{k+1} - 16^N) \\ &= \left((4^{2N+2} - 40) \cdot 4^k + 50 \cdot 16^N - 32 \cdot 256^N - 4 \cdot 64^N \right) 4^k + (8 \cdot 1024^N - 10 \cdot 256^N) \\ &> 0. \end{aligned} \quad (A.1)$$

³The inequality (3.28) is proved in the appendix.

We see that for $k \geq 2N + 1$,

$$\begin{aligned} & (4^{2N+2} - 40) \cdot 4^k + 50 \cdot 16^N - 32 \cdot 256^N - 4 \cdot 64^N \\ & > (4^{2N+2} - 40) \cdot 4^{2N+1} + 50 \cdot 16^N - 32 \cdot 256^N - 4 \cdot 64^N \\ & = 224 \cdot 256^N - 590 \cdot 16^N - 4 \cdot 64^N > 0 \end{aligned}$$

and

$$8 \cdot 1024^N - 10 \cdot 256^N > 0.$$

Hence, (A.1) holds for $k \geq 2N + 1$.

Appendix B: Proof of (3.21)

$$\begin{aligned} q_n - 20p_n &= (3780n^3 - 22680n^2 - 112455n + 273105)4^n + 128n^9 - 576n^8 + 5248n^7 \\ &\quad + 2016n^6 - 32984n^5 + 70476n^4 + 250052n^3 - 134916n^2 - 512244n - 226800 \\ &= (179550 + 397845(n-9) + 79380(n-9)^2 + 3780(n-9)^3)4^n \\ &\quad + 49648561200 + 46968464520(n-9) + 19975332000(n-9)^2 \\ &\quad + 5019956996(n-9)^3 + 822741108(n-9)^4 + 91303912(n-9)^5 \\ &\quad + 6864480(n-9)^6 + 337024(n-9)^7 + 9792(n-9)^8 + 128(n-9)^9 \\ &> 0 \quad \text{for } n \geq 9. \end{aligned}$$

Appendix C: Proof of (3.28)

We now show that for $n \geq 6$,

$$\begin{aligned} Q_n - 12P_n &= (16n^7 - 56n^6 + 268n^5 - 70412n^3 + 252112n^2 + 909099n - 1556415)9^n \\ &\quad + 70n^4 \cdot 9^n - (68040n^4 + 476280n^3 + 413910n^2 - 1173690n - 1179360)4^n \\ &\quad - (11664n^7 - 40824n^6 + 39852n^5 + 595350n^4 + 256932n^3 - 485352n^2 \\ &\quad - 106029n - 104895) > 0. \end{aligned}$$

It suffices to show that for $n \geq 6$,

$$\left(\frac{9}{4}\right)^n > A_n, \tag{C.1}$$

where

$$A_n = \frac{68040n^4 + 476280n^3 + 413910n^2 - 1173690n - 1179360}{16n^7 - 56n^6 + 268n^5 - 70412n^3 + 252112n^2 + 909099n - 1556415},$$

and

$$\begin{aligned} \cdot 9^n &> \frac{1}{70n^4} \left(11664n^7 - 40824n^6 + 39852n^5 + 595350n^4 + 256932n^3 - 485352n^2 \right. \\ &\quad \left. - 106029n - 104895 \right). \end{aligned} \tag{C.2}$$

By induction with respect to n , we can prove the inequalities (C.1) and (C.2). Here, we only prove the inequality (C.1). The proof of (C.2) is analogous.

For $n = 6$ in (C.1), we find that

$$\left(\frac{9}{4}\right)^6 = \frac{531441}{4096} = 129.746\dots \quad \text{and} \quad A_6 = \frac{3138660}{27229} = 115.269\dots$$

This shows that (C.1) holds for $n = 6$.

Now we assume that (C.1) holds for some $n \geq 6$. Then, for $n \mapsto n + 1$ in (C.1), by using the induction hypothesis, we have

$$\left(\frac{9}{4}\right)^{n+1} - A_{n+1} > \frac{9}{4}A_n - A_{n+1} = \frac{2835R_n}{2S_nT_n},$$

where

$$\begin{aligned} R_n &= 960n^{11} + 12384n^{10} + 73088n^9 + 256200n^8 - 3508908n^7 - 22121984n^6 + 50474996n^5 \\ &\quad + 274552068n^4 - 445858781n^3 - 777353865n^2 + 997107984n - 660306024 \\ &= 878926761468 + 1894841991720(n-6) + 1695853296525(n-6)^2 \\ &\quad + 849645117283(n-6)^3 + 268187103036(n-6)^4 + 56595283460(n-6)^5 \\ &\quad + 8234103112(n-6)^6 + 835076820(n-6)^7 + 58479432(n-6)^8 + 2716928(n-6)^9 \\ &\quad + 75744(n-6)^{10} + 960(n-6)^{11}, \end{aligned}$$

$$\begin{aligned} S_n &= 16n^7 - 56n^6 + 268n^5 - 70412n^3 + 252112n^2 + 909099n - 1556415 \\ &= 1715427 + 679323(n-6) + 1087672(n-6)^2 + 509908(n-6)^3 + 98760(n-6)^4 \\ &\quad + 10348(n-6)^5 + 616(n-6)^6 + 16(n-6)^7 \end{aligned}$$

and

$$\begin{aligned} T_n &= 16n^7 + 56n^6 + 268n^5 + 1060n^4 - 68292n^3 + 43052n^2 + 1203203n - 465388 \\ &= 4102070 + 4834979(n-6) + 3323012(n-6)^2 + 1021308(n-6)^3 + 160300(n-6)^4 \\ &\quad + 14380(n-6)^5 + 728(n-6)^6 + 16(n-6)^7. \end{aligned}$$

Hence, we have

$$\left(\frac{9}{4}\right)^{n+1} > A_{n+1}.$$

Thus, by the principle of mathematical induction, the inequality (C.1) holds for $n \geq 6$.

ACKNOWLEDGEMENT

Some computations in this paper were performed using Maple software.

REFERENCES

- [1] A. Baricz, J. Sándor, Extensions of generalized Wilker inequality to Bessel functions, *J. Math. Inequal.* 2 (2008) 397–406.
- [2] C.-P. Chen, W.-S. Cheung, Inequalities and solution to Oppenheim's problem, *Integral Transforms Spec. Funct.* 23 (2012), no. 5, 325–336.
- [3] C.-P. Chen, W.-S. Cheung, Sharpness of Wilker and Huygens type inequalities, *J. Inequal. Appl.* 2012 (2012) 72, <http://dx.doi.org/10.1186/1029-242X-2012-72>.
- [4] C.-P. Chen, R.B. Paris, Series representations of the remainders in the expansions for certain trigonometric and hyperbolic functions with applications, <http://arxiv.org/abs/1601.03180>.
- [5] C.-P. Chen, F. Qi, A double inequality for remainder of power series of tangent function, *Tamkang J. Math.* 34 (2003), no.4, 351–355.
- [6] C.-P. Chen, J. Sándor, Inequality chains for Wilker, Huygens and Lazarević type inequalities, *J. Math. Inequal.* 8 (2014), no. 1, 55–67.
- [7] C. D'aniello, On some inequalities for the Bernoulli numbers, *Rend. Circ. Mat. Palermo* 43 (1994) 329–332.

- [8] B.-N. Guo, B.-M. Qiao, F. Qi, W. Li, On new proofs of Wilker inequalities involving trigonometric functions, *Math. Inequal. Appl.* 6 (2003) 19–22.
- [9] C. Huygens, *Oeuvres Completes 1888-1940*, Société Hollandaise des Science, Haga.
- [10] C. Mortici, The natural approach of Wilker-Cusa-Huygens inequalities, *Math. Inequal. Appl.* 14 (2011) 535–541.
- [11] C. Mortici, A subtly analysis of Wilker inequality, *Appl. Math. Comput.* 231 (2014) 516–520.
- [12] E. Neuman, One- and two-sided inequalities for Jacobian elliptic functions and related results, *Integral Transforms Spec. Funct.* 21 (2010) 399–407.
- [13] E. Neuman, On Wilker and Huygens type inequalities, *Math. Inequal. Appl.* 15 (2012), no. 2, 271–279.
- [14] E. Neuman, Wilker and Huygens-type inequalities for the generalized trigonometric and for the generalized hyperbolic functions, *Appl. Math. Comput.* 230 (2014) 211–217.
- [15] E. Neuman, J. Sándor, On some inequalities involving trigonometric and hyperbolic functions with emphasis on the Cusa-Huygens, Wilker, and Huygens inequalities, *Math. Inequal. Appl.* 13 (2010) 715–723.
- [16] I. Pinelis, L'Hospital rules of monotonicity and Wilker-Anglesio inequality, *Amer. Math. Monthly*, 111 (2004) 905–909.
- [17] J. S. Sumner, A. A. Jagers, M. Vowe, J. Anglesio, Inequalities involving trigonometric functions, *Amer. Math. Monthly* 98 (1991) 264–267.
- [18] J. B. Wilker, Problem E 3306, *Amer. Math. Monthly* 96 (1989) 55.
- [19] S.-H. Wu, On extension and refinement of Wilker's inequality, *Rocky Mountain J. Math.* 39 (2009) 683–687.
- [20] S.-H. Wu, A. Baricz, Generalizations of Mitrinović, Adamović and Lazarević's inequalities and their applications, *Publ. Math. Debrecen* 75 (2009) 447–458.
- [21] S.-H. Wu, H. M. Srivastava, A weighted and exponential generalization of Wilker's inequality and its applications, *Integral Transforms and Spec. Funct.* 18 (2007) 529–535.
- [22] S.-H. Wu and H. M. Srivastava, A further refinement of Wilker's inequality, *Integral Transforms and Spec. Funct.* 19 (2008) 757–765.
- [23] L. Zhang and L. Zhu, A new elementary proof of Wilker's inequalities, *Math. Inequal. Appl.* 11 (2008) 149–151.
- [24] J.-L. Zhao, Q.-M. Luo, B.-N. Guo, F. Qi, Remarks on inequalities for the tangent function. *Hacet. J. Math. Stat.* 41 (2012), no. 4, 499–506.
- [25] L. Zhu, A new simple proof of Wilker's inequality, *Math. Inequal. Appl.* 8 (2005) 749–750.
- [26] L. Zhu, Some new Wilker-type inequalities for circular and hyperbolic functions, *Abstr. Appl. Anal.* 2009 (2009), Article ID 485842, <http://www.hindawi.com/journals/aaa/2009/485842/>.
- [27] L. Zhu, A source of inequalities for circular functions, *Comput. Math. Appl.* 58 (2009) 1998–2004.
- [28] L. Zhu, Inequalities for Hyperbolic functions and their Applications, *J. Inequal. Appl.* 2010 (2010), Article ID 130821, <http://www.emis.de/journals/HOA/JIA/Volume2010/130821.abs.html>.

C.-P. CHEN: SCHOOL OF MATHEMATICS AND INFORMATICS, HENAN POLYTECHNIC UNIVERSITY, JIAOZUO CITY 454000, HENAN PROVINCE, CHINA

E-mail address: chenchaoping@sohu.com

R.B. PARIS: DIVISION OF COMPUTING AND MATHEMATICS,, UNIVERSITY OF ABERTAY, DUNDEE, DD1 1HG, UK

E-mail address: R.Paris@abertay.ac.uk